

Presentation Type: Poster

Track: Aquatic Toxicology and Ecology

Session: TBD

Abstract Title: Development and initial evaluation of a reconstituted water formulation that better represents natural waters.

Authors: S.M. Hoheisel, c/o U.S. EPA, Duluth, MN
R.J. Erickson, T.L. Highland, J.R. Hockett, D.J. Hoff T.J. Norberg-King, U.S. EPA,
Duluth, MN
T.R. Valenti, National Research Council, c/o U.S. EPA, Duluth, MN
D.R. Mount, U.S. EPA, Duluth, MN

Abstract:

The use of reconstituted waters is deeply entrenched in many standardized aquatic toxicity testing protocols. The primary appeal of reconstituted waters is inter-laboratory comparability, such that experiments performed in different laboratories can be conducted in (nominally) identical waters. While several formulations of reconstituted waters have been developed over the years, many aquatic toxicity test protocols from the U.S. EPA and ASTM use a formulation that has its roots in a publication by Marking and Dawson (1973). While widely used, the ionic composition of these waters is not typical of North American surface waters; for example, the molar Ca:Mg ratio in EPA/ASTM water is 0.70, but is averages about 2.2 in U.S. waters. More dramatic are the differences in molar Cl:SO₄ ratio, which is 0.0632 in EPA/ASTM water, but is more typically in the range of 1 to 2 in U.S. waters., with characteristics such as too much Mg relative to Ca, and too much sulfate relative to chloride. We have developed a new reconstituted water formulation based on an analysis of natural waters from across the U.S. Hardness was chosen as the master variable from which the remaining composition is calculated. The subsequent relationships that define the water composition are alkalinity v. hardness, Ca:Mg ratio, Cl:SO₄ chloride:sulfate ratio v. hardness, and K as a proportion of Na + K. Exactly mimicking natural waters is tedious in the laboratory, because natural waters gain alkalinity from calcium and magnesium carbonates, salts which are difficult to dissolve; this formulation uses only readily soluble salts, though this logistical convenience sacrifices some fidelity to the central tendency of natural waters. Recipes for water of any hardness (recommended bounds of 10 to 400 mg/L as calcium carbonate) can be calculated in a simple spreadsheet. Initial evaluation of these waters in chronic tests with *Ceriodaphnia dubia* and *Daphnia magna*, and short-term chronic tests with fathead minnows (*Pimephales promelas*), have shown performance in our proposed reconstituted-the-reformulated water to compare favorably with Marking and Dawson-based waters or natural Lake Superior water. Additional testing with *Chironomus dilutus* and *Hyalella azteca* is planned. Using reconstituted waters that are more representative of natural waters should improve assessment of chemicals whose toxicity is influenced by the ionic composition of the water (e.g., many cationic metals), and may also improve performance of some organisms found to be sensitive to the ionic composition of water (e.g., *Hyalella*).